

# Results

	Bariatric Surgery (n = 55,789)	Nonsurgical Control (n = 55,789)	Hazard Ratio	95% Confidence Interval	p-value
Obesity-associated cancers	2,206 (4.0%)	4,960 (8.9%)	0.482	0.459-0.507	0.000
Esophageal adenocarcinoma	10 (<0.1%)	10 (<0.1%)	0.819	0.333-2.011	0.242
Breast	501 (0.9%)	751 (1.3%)	0.753	0.678-0.836	0.001
Renal	104 (0.2%)	132 (0.2%)	0.816	0.646-1.031	0.744
Colon	201 (0.4%)	360 (0.6%)	0.638	0.541-0.752	0.000
Rectal	29 (0.1%)	50 (0.1%)	0.670	0.423-1.060	0.561
Gastric	11 (<0.1%)	12 (<0.1%)	1.137	0.517-2.502	0.126
Liver	969 (1.7%)	2,198 (4.1%)	0.370	0.345-0.396	0.000
Gallbladder	10 (<0.1%)	10 (<0.1%)	0.390	0.125-1.213	0.372
Pancreas	54 (0.1%)	86 (<0.1%)	0.784	0.569-1.080	0.028
Ovarian	130 (0.2%)	214 (0.4%)	0.654	0.531-0.806	0.002
Endometrial	105 (0.2%)	245 (0.4%)	0.448	0.362-0.556	0.979
Thyroid	154 (0.3%)	175 (0.3%)	0.873	0.715-1.065	0.000
Multiple Myeloma	44 (0.1%)	64 (0.1%)	0.753	0.519-1.09	0.021